Lochs' index: weight function and change of basis

Pablo Rotondo

LIGM, Université Gustave Eiffel

Joint work with Valérie Berthé, Eda Cesaratto and Martín Safe

Meeting STIC-AmSud, RAPA2, Online, 7 December, 2021.

Motivation: simulating continued fractions

For computer simulation:

• Given t binary digits b_1, b_2, \ldots, b_t of $x \in [0, 1]$,

$$x = (0.b_1b_2\ldots)_2 \in [0,1].$$

Number $n = n_t(x)$ of CFE-digits (partial quotients) deduced without possibility of error ?

$$x = \frac{1}{a_1 + \frac{1}{a_2 + \cdots}}$$

Motivation: simulating continued fractions

For computer simulation:

• Given t binary digits b_1, b_2, \ldots, b_t of $x \in [0, 1]$,

$$x = (0.b_1b_2\ldots)_2 \in [0,1].$$

Number $n = n_t(x)$ of CFE-digits (partial quotients) deduced without possibility of error ?

$$x = \frac{1}{a_1 + \frac{1}{a_2 + \cdots}}$$

Natural to consider the quotient $n_t(x)/t$:

- rate of CFE digits per binary digit,
- compares relative information/redundancy of expansions.

First historical results: Lochs' Theorem

Theorem (Lochs '64)

The rate of CF-digits per decimal given satisfies

$$\lim_{\mathbf{d} \to \infty} \frac{n_{\mathbf{d}}(x)}{\mathbf{d}} = \frac{6 \log 2 \log 10}{\pi^2} \doteq 0.9702701 \dots \,,$$

for almost every x.

First historical results: Lochs' Theorem

Theorem (Lochs '64)

The rate of CF-digits per decimal given satisfies

$$\lim_{\mathbf{d} \to \infty} \frac{n_{\mathbf{d}}(x)}{\mathbf{d}} = \frac{6 \log 2 \log 10}{\pi^2} \doteq 0.9702701 \dots \,,$$

for almost every x.

"Example". The first 1000 decimals of π determine exactly 968 partial quotients of π .

First historical results: Lochs' Theorem

Theorem (Lochs '64)

The rate of CF-digits per decimal given satisfies

$$\lim_{\mathbf{d} \to \infty} \frac{n_{\mathbf{d}}(x)}{\mathbf{d}} = \frac{6 \log 2 \log 10}{\pi^2} \doteq 0.9702701 \dots \,,$$

for almost every x.

"Example". The first 1000 decimals of π determine exactly 968 partial quotients of π .

Theorem (Faivre '98)

$$\Pr\left\{x\in[0,1]:\frac{n_{\mathsf{d}}(x)-\mathsf{d}\times a}{\sigma\sqrt{\mathsf{d}}}\leq\theta\right\}\to \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\theta}e^{-u^2/2}du\,,$$

where $a = \frac{6 \log 2 \log 10}{\pi^2}$ is the Lochs' constant, and $\sigma > 0$.

• Given t binary digits $b_1, b_2, \ldots, b_t \in \{0, 1\}$ of

$$x = (0.b_1b_2\ldots)_2 \in [0,1].$$

Number $n = n_t(x)$ of *d*-ary digits $0 \le d_1, \ldots, d_n < d$ deduced?

$$x = (0.d_1d_2...)_d \in [0,1].$$

• Given t binary digits $b_1, b_2, \ldots, b_t \in \{0, 1\}$ of

$$x = (0.b_1b_2\ldots)_2 \in [0,1].$$

Number $n = n_t(x)$ of *d*-ary digits $0 \le d_1, \ldots, d_n < d$ deduced?

$$x = (0.d_1d_2...)_d \in [0,1].$$

Answer:

• For $d = 2^A$ we simply obtain

$$A \times n_t(x) = t \,,$$

because 1 d-ary digit corresponds to A binary digits.

• Given t binary digits $b_1, b_2, \ldots, b_t \in \{0, 1\}$ of

$$x = (0.b_1b_2\ldots)_2 \in [0,1].$$

Number $n = n_t(x)$ of *d*-ary digits $0 \le d_1, \ldots, d_n < d$ deduced?

$$x = (0.d_1d_2...)_d \in [0,1].$$

Answer:

• For $d = 2^A$ we simply obtain

$$A \times n_t(x) = t \,,$$

because 1 d-ary digit corresponds to A binary digits.

More generally we expect

$$\frac{\log d}{\log 2} \times n_t(x) \sim t \,,$$

• Given t binary digits $b_1, b_2, \ldots, b_t \in \{0, 1\}$ of

$$x = (0.b_1b_2\ldots)_2 \in [0,1].$$

Number $n = n_t(x)$ of *d*-ary digits $0 \le d_1, \ldots, d_n < d$ deduced?

$$x = (0.d_1d_2...)_d \in [0,1].$$

Answer:

• For $d = 2^A$ we simply obtain

$$A \times n_t(x) = t \,,$$

because 1 d-ary digit corresponds to A binary digits.

More generally we expect

$$\frac{\log d}{\log 2} \times n_t(x) \sim t \,,$$

one digit in base d^B corresponds to one in base 2^A when $d^B \approx 2^A$.

Classical

▶ Dajani&Fieldsteel'01: From source S₁ to S₂, both of positive entropy:

 $\lim L_t(x; S_1, S_2)/t = h(S_1)/h(S_2),$

where $L_t(x; S_1, S_2)$ is number of digits in S_2 deduced from t in S_1 .

Classical

▶ Dajani&Fieldsteel'01: From source S₁ to S₂, both of positive entropy:

 $\lim L_t(x; S_1, S_2)/t = h(S_1)/h(S_2),$

where $L_t(x; S_1, S_2)$ is number of digits in S_2 deduced from t in S_1 .

Classical

▶ Dajani&Fieldsteel'01: From source S₁ to S₂, both of positive entropy:

$$\lim L_t(x; S_1, S_2)/t = h(S_1)/h(S_2),$$

where $L_t(x; S_1, S_2)$ is number of digits in S_2 deduced from t in S_1 .

• What if
$$h(S_1) = 0$$
 or $h(S_2) = 0$?

- If $h(S_2) = 0$ and $h(S_1) > 0$, almost surely $L/t \to \infty$.

– If $h(S_2) > 0$ and $h(S_1) = 0$, almost surely $L/t \to 0$.

Classical

▶ Dajani&Fieldsteel'01: From source S₁ to S₂, both of positive entropy:

$$\lim L_t(x; S_1, S_2)/t = h(S_1)/h(S_2),$$

where $L_t(x; S_1, S_2)$ is number of digits in S_2 deduced from t in S_1 .

• What if
$$h(S_1) = 0$$
 or $h(S_2) = 0$?

- If
$$h(S_2) = 0$$
 and $h(S_1) > 0$, almost surely $L/t \to \infty$.

- If
$$h(S_2) > 0$$
 and $h(S_1) = 0$, almost surely $L/t \to 0$.

Our work

- ▶ Introduce appropriate notion of *renormalized* entropy f_1, f_2 ,
- Generalization: for positive, zero or infinite entropy:

$$\lim f_2(L_t(x; S_1, S_2))/f_1(t) = 1.$$

Plan of the talk

1. Definitions: partitions, Lochs' and weight function

- 2. Statement of main result and discussion
- 3. Examples of natural zero entropy sources that have weight
- 4. Concepts for the proof of the main result
- 5. Conclusions



1. Definitions: partitions, Lochs' and weight function

2. Statement of main result and discussion

3. Examples of natural zero entropy sources that have weight

4. Concepts for the proof of the main result

5. Conclusions

Intervals: sources and partitions

Definition (System of interval partitions)

Sequence of topological partitions $\mathcal{P} = (\mathcal{P}_n)$ of [0,1]

•
$$\mathcal{P}_{n+1}$$
 refinement of \mathcal{P}_n for every n .

▶
$$\|\mathcal{P}_n\| = \sup\{\mathtt{diam}(I) : I \in \mathcal{P}_n\}$$
 tends to 0.

Intervals: sources and partitions

Definition (System of interval partitions)

Sequence of topological partitions $\mathcal{P} = (\mathcal{P}_n)$ of [0,1]

•
$$\mathcal{P}_{n+1}$$
 refinement of \mathcal{P}_n for every n .

▶
$$\|\mathcal{P}_n\| = \sup\{\mathtt{diam}(I) : I \in \mathcal{P}_n\}$$
 tends to 0.

Equivalent to sources

• notation
$$I_n^{\mathcal{P}}(x) = I \in \mathcal{P}_n$$
 such that $x \in I$,

• first *n* symbols for *x* determine $I_n^{\mathcal{P}}(x)$ and conversely.

Intervals: sources and partitions

Definition (System of interval partitions)

Sequence of topological partitions $\mathcal{P} = (\mathcal{P}_n)$ of [0,1]

•
$$\mathcal{P}_{n+1}$$
 refinement of \mathcal{P}_n for every n .

▶
$$\|\mathcal{P}_n\| = \sup\{\mathtt{diam}(I) : I \in \mathcal{P}_n\}$$
 tends to 0.

Equivalent to sources

• notation
$$I_n^{\mathcal{P}}(x) = I \in \mathcal{P}_n$$
 such that $x \in I$,

• first *n* symbols for *x* determine $I_n^{\mathcal{P}}(x)$ and conversely.

Example. Decimal expansion

Depth n interval for $x = (0.d_1d_2...)_{10}$

$$I_n^{\mathcal{D}}(x) = \left((0.d_1 \dots d_n)_{10}, (0.d_1 \dots d_n)_{10} + 10^{-n} \right) + 0^{-n}$$

containing $y \in (0,1)$ having the exact same first n digits as x.

Entropy of a partition

Entropy dictates size of intervals

► Shannon entropy¹:

$$H(\mathcal{P}) = -\lim_{k \to \infty} \frac{1}{k} \sum_{I \in \mathcal{P}_k} |I| \log |I| .$$

 \blacktriangleright *Point-wise*: for almost every x

$$h(\mathcal{P}) = -\lim_{k \to \infty} \frac{1}{k} \log \left| I_k^{\mathcal{P}}(x) \right| \,.$$

¹We consider Lebesgue measure here, but any Borel λ works.

Entropy of a partition

Entropy dictates size of intervals

► Shannon entropy¹:

$$H(\mathcal{P}) = -\lim_{k \to \infty} \frac{1}{k} \sum_{I \in \mathcal{P}_k} |I| \log |I| .$$

• *Point-wise*: for almost every x

$$h(\mathcal{P}) = -\lim_{k \to \infty} \frac{1}{k} \log \left| I_k^{\mathcal{P}}(x) \right| \,.$$

Point-wise to Shannon

$$H(\mathcal{P}) = \lim_{k \to \infty} \mathbb{E}\left[-\frac{1}{k} \log \left|I_k^{\mathcal{P}}(x)\right|\right].$$

¹We consider Lebesgue measure here, but any Borel λ works.

Generalization Lochs': Lochs' index

The Lochs' index

- formalizes the notation of deduced digits,
- generalizes it to systems of interval partitions.

Generalization Lochs': Lochs' index

The Lochs' index

- formalizes the notation of deduced digits,
- generalizes it to systems of interval partitions.

Lochs' index for systems of partitions $\mathcal{P}^1, \mathcal{P}^2$

$$L_n(x; \mathcal{P}^1, \mathcal{P}^2) := \max\{m \ge 0 : I_n^{\mathcal{P}^1}(x) \subset I_m^{\mathcal{P}^2}(x)\},\$$

depth in \mathcal{P}^2 deduced from depth n in \mathcal{P}^1 .

Generalization Lochs': Lochs' index

The Lochs' index

- formalizes the notation of deduced digits,
- generalizes it to systems of interval partitions.

Lochs' index for systems of partitions $\mathcal{P}^1, \mathcal{P}^2$

$$L_n(x; \mathcal{P}^1, \mathcal{P}^2) := \max\{m \ge 0 : I_n^{\mathcal{P}^1}(x) \subset I_m^{\mathcal{P}^2}(x)\},\$$

depth in \mathcal{P}^2 deduced from depth n in \mathcal{P}^1 .

Explanation

If $I_n^{\mathcal{P}^1}(x)$ splits over (intersects) several $J \in \mathcal{P}_m^2$, \implies we cannot yet decide on $I_m^{\mathcal{P}^2}(x)$

Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with positive point-wise entropies $h(\mathcal{P}^1)$ and $h(\mathcal{P}^2)$. Then

$$\lim_{n \to \infty} \frac{1}{n} L_n(x; \mathcal{P}^1, \mathcal{P}^2) = \frac{h(\mathcal{P}^1)}{h(\mathcal{P}^2)}$$

for a.e. x.

Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with positive point-wise entropies $h(\mathcal{P}^1)$ and $h(\mathcal{P}^2)$. Then

$$\lim_{n \to \infty} \frac{1}{n} L_n(x; \mathcal{P}^1, \mathcal{P}^2) = \frac{h(\mathcal{P}^1)}{h(\mathcal{P}^2)}$$

for a.e. x.

– Base d. Since $|I_t^{\mathcal{D}}(x)| = d^{-t}$, $h(\mathcal{D}) = \log d$.

Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with positive point-wise entropies $h(\mathcal{P}^1)$ and $h(\mathcal{P}^2)$. Then

$$\lim_{n \to \infty} \frac{1}{n} L_n(x; \mathcal{P}^1, \mathcal{P}^2) = \frac{h(\mathcal{P}^1)}{h(\mathcal{P}^2)}$$

for a.e. x.

- Base d. Since $|I_t^{\mathcal{D}}(x)| = d^{-t}$, $h(\mathcal{D}) = \log d$.
- Continued fractions. Intervals satisfy $|I_k^{\mathcal{C}}(x)| = \Theta((q_k(x))^{-2})$

$$h(\mathcal{C}) = 2\lim_{k \to \infty} \frac{1}{k} \log q_k(x) = \frac{\pi^2}{6 \log 2}$$

Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with positive point-wise entropies $h(\mathcal{P}^1)$ and $h(\mathcal{P}^2)$. Then

$$\lim_{n \to \infty} \frac{1}{n} L_n(x; \mathcal{P}^1, \mathcal{P}^2) = \frac{h(\mathcal{P}^1)}{h(\mathcal{P}^2)}$$

for a.e. x.

- Base d. Since $|I_t^{\mathcal{D}}(x)| = d^{-t}$, $h(\mathcal{D}) = \log d$.
- Continued fractions. Intervals satisfy $|I_k^{\mathcal{C}}(x)| = \Theta((q_k(x))^{-2})$

$$h(\mathcal{C}) = 2\lim_{k \to \infty} \frac{1}{k} \log q_k(x) = \frac{\pi^2}{6 \log 2}$$

 \implies we deduce Lochs' Theorem and the result for d-ary basis.

Existence of point-wise entropy

Systems of partitions associated with good (positive entropy) dynamical systems have point-wise entropy:

Theorem (Shannon, McMillan, Breiman)

Let T be an ergodic measure preserving transformation on a probability space $(\Omega, \mathcal{B}, \mu)$ and let P be a finite or countable generating partition for T for which $H_{\mu}(P) < \infty$. Then for μ -a.e. x,

$$\lim_{n \to \infty} -\frac{\log \mu \left(P_n(x) \right)}{n} = h_\mu(T) \,.$$

Here $H_{\mu}(P)$ denotes the entropy of the partition P, $h_{\mu}(T)$ the entropy of T and $P_n(x)$ denotes the element of the partition $\bigvee_{i=0}^{n-1} T^{-i}P$ containing x.

Log-balancedness and weight function

Definition (Weight function)

A system of partitions $\mathcal{P} = (\mathcal{P}_n)$ is *log-balanced* a.e. (resp. in measure) with *weight function* $f \colon \mathbb{N} \to \mathbb{R}_{>0}$, $f(n) \to \infty$, if

$$-\log|I_n^{\mathcal{P}}(x)| \sim f(n) \,,$$

almost everywhere (resp. in measure).

Log-balancedness and weight function

Definition (Weight function)

A system of partitions $\mathcal{P} = (\mathcal{P}_n)$ is *log-balanced* a.e. (resp. in measure) with *weight function* $f \colon \mathbb{N} \to \mathbb{R}_{>0}$, $f(n) \to \infty$, if

$$-\log|I_n^{\mathcal{P}}(x)| \sim f(n),$$

almost everywhere (resp. in measure).

Example

For positive entropy
$$h = h(\mathcal{P}) > 0$$

$$f(n) = h \times n \,.$$

▶ If partition is log-balanced, entropy 0 corresponds to

$$f(n) = o(n) \, .$$

Realization result for weight functions

Proposition

Let $f: \mathbb{N} \to \mathbb{R}_{>0}$ non-decreasing, $f(n) \to \infty$. Then there exists a log-balanced \mathcal{P} with weight function f almost everywhere.

Realization result for weight functions

Proposition

Let $f: \mathbb{N} \to \mathbb{R}_{>0}$ non-decreasing, $f(n) \to \infty$. Then there exists a log-balanced \mathcal{P} with weight function f almost everywhere.

Proof sketch.

Given n, let k=k(n) be such that $2^k \leq \exp(f(n)) < 2^{k+1}.$ Define

$$\mathcal{P}_n := \left\{ \left(\frac{i}{2^k}, \frac{i+1}{2^k}\right) : 0 \le i < 2^k \right\},$$

so that $|I_n(x)| = 2^{-k}$ satisfies $e^{-f(n)} \le 2^{-k} < 2e^{-f(n)}$.



1. Definitions: partitions, Lochs' and weight function

2. Statement of main result and discussion

3. Examples of natural zero entropy sources that have weight

4. Concepts for the proof of the main result

5. Conclusions

Our main result

Theorem (Berthé, Cesaratto, R., Safe, 2021+) Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with a.e. weight functions f_1 and f_2 . Then, under certain technical conditions

$$\lim_{n \to \infty} \frac{f_2\left(L_n(x; \mathcal{P}^1, \mathcal{P}^2)\right)}{f_1(n)} = 1,$$

for a.e. x.

Our main result

Theorem (Berthé, Cesaratto, R., Safe, 2021+) Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with a.e. weight functions f_1 and f_2 . Then, under certain technical conditions

$$\lim_{n \to \infty} \frac{f_2\left(L_n(x; \mathcal{P}^1, \mathcal{P}^2)\right)}{f_1(n)} = 1,$$

for a.e. x.

The conditions are:

•
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

• f_2 is non decreasing ;

▶
$$f_2(n+1) - f_2(n) = o(f_2(n))$$
 as $n \to \infty$.
Our main result

Theorem (Berthé, Cesaratto, R., Safe, 2021+) Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with a.e. weight functions f_1 and f_2 . Then, under certain technical conditions

$$\lim_{n \to \infty} \frac{f_2\left(L_n(x; \mathcal{P}^1, \mathcal{P}^2)\right)}{f_1(n)} = 1,$$

for a.e. x.

The conditions are:

•
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

• f_2 is non decreasing ;

▶
$$f_2(n+1) - f_2(n) = o(f_2(n))$$
 as $n \to \infty$.

Remark. First condition can be dropped for convergence in measure.

$$(a) \ \sum e^{-\delta f_1(n)} < \infty \ \text{for every} \ \delta > 0;$$

(b) f_2 is non decreasing ;

(c)
$$f_2(n+1) - f_2(n) = o(f_2(n))$$
 as $n \to \infty$.

(a)
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

(b) f_2 is non decreasing ;

 $(c) \ \ f_2(n+1) - f_2(n) = o(f_2(n)) \text{ as } n \to \infty.$

Intuitively, the first condition is the most constraining one:

- Condition (b) reflects the fact that \mathcal{P}_2 is refining ;
- Condition (c) means that $f_2(n+1) \sim f_2(n)$;
- Condition (a) tells us that $f_1(n)$ grows not too slowly

(a)
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

(b) f_2 is non decreasing ;

 $(c) \ f_2(n+1) - f_2(n) = o(f_2(n)) \text{ as } n \to \infty.$

Intuitively, the first condition is the most constraining one:

- Condition (b) reflects the fact that \mathcal{P}_2 is refining ;
- Condition (c) means that $f_2(n+1) \sim f_2(n)$;
- Condition (a) tells us that $f_1(n)$ grows not too slowly

Important remarks

- Condition (a) not satisfied when $f_1(n) = \log n$,
- Condition (a) satisfied for $f_1(n) \ge (\log n)^2$.

(a)
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

(b) f_2 is non decreasing ;

 $(c) \ f_2(n+1) - f_2(n) = o(f_2(n)) \text{ as } n \to \infty.$

Intuitively, the first condition is the most constraining one:

- Condition (b) reflects the fact that \mathcal{P}_2 is refining ;
- Condition (c) means that $f_2(n+1) \sim f_2(n)$;
- Condition (a) tells us that $f_1(n)$ grows not too slowly

Important remarks

- Condition (a) not satisfied when $f_1(n) = \log n$,
- Condition (a) satisfied for $f_1(n) \ge (\log n)^2$.
- Condition (c) not satisfied when $f_2(n) = \exp(n)$,
- Condition (c) is satisfied when $f_2(n) = \exp(\sqrt{n})$.

Discussion: conditions of our main result

Example: appropriate output partitions \mathcal{P}_2 Subexponential weight functions of the form

 $f_2(n) = \exp(g(n))\,,$

with $g'(t) \searrow 0$.

Discussion: conditions of our main result

Example: appropriate output partitions \mathcal{P}_2 Subexponential weight functions of the form

 $f_2(n) = \exp(g(n))\,,$

with $g'(t) \searrow 0$.

Example: appropriate input partitions \mathcal{P}_1

Superlogarithmic weight functions

 $f_1(n) = (\log n) \cdot g(n) \,,$

with $g(t) \to \infty$.

Discussion: conditions of our main result

Example: appropriate output partitions \mathcal{P}_2 Subexponential weight functions of the form

 $f_2(n) = \exp(g(n))\,,$

with $g'(t) \searrow 0$.

Example: appropriate input partitions \mathcal{P}_1

Superlogarithmic weight functions

 $f_1(n) = (\log n) \cdot g(n) \,,$

with $g(t) \to \infty$.

Note. For convergence in measure the conditions on the input partitions can be dropped.



1. Definitions: partitions, Lochs' and weight function

2. Statement of main result and discussion

3. Examples of natural zero entropy sources that have weight

4. Concepts for the proof of the main result

5. Conclusions

Farey partition (Sturm source) and Stern-Brocot partition built by splitting intervals at *mediant*

$$\operatorname{mediant}(a/b,c/d) := (a+b)/(c+d)$$
.

Farey partition (Sturm source) and Stern-Brocot partition built by splitting intervals at *mediant*

$$\texttt{mediant}(a/b,c/d) := (a+b)/(c+d)$$
.

Farey partition \mathcal{F}_n :

- Base case: $\mathcal{F}_0 = \{[0,1]\}.$
- ▶ Building \mathcal{F}_n : split $\begin{bmatrix} a \\ b \end{bmatrix}$, $\begin{bmatrix} c \\ d \end{bmatrix}$ ∈ \mathcal{F}_{n-1} , if $b + d \le n + 1$.

Stern-Brocot partition SB_n :

- Base case: $SB_0 = \{[0, 1]\}.$
- ▶ Building SB_n : split $\left[\frac{a}{b}, \frac{c}{d}\right] \in SB_{n-1}$ always.

Farey partition (Sturm source) and Stern-Brocot partition built by splitting intervals at *mediant*

$$\operatorname{mediant}(a/b,c/d) := (a+b)/(c+d)$$
.

Farey partition \mathcal{F}_n :

- Base case: $\mathcal{F}_0 = \{[0,1]\}.$
- ▶ Building \mathcal{F}_n : split $\begin{bmatrix} a \\ b \end{bmatrix}$, $\begin{bmatrix} c \\ d \end{bmatrix}$ ∈ \mathcal{F}_{n-1} , if $b + d \le n + 1$.

Stern-Brocot partition SB_n :

- Base case: $SB_0 = \{[0, 1]\}.$
- ▶ Building SB_n : split $\begin{bmatrix} \frac{a}{b}, \frac{c}{d} \end{bmatrix} \in SB_{n-1}$ always.



Farey partition (Sturm source) and Stern-Brocot partition built by splitting intervals at *mediant*

$$\texttt{mediant}(a/b,c/d) := (a+b)/(c+d)$$
.

Farey partition \mathcal{F}_n :

• Base case:
$$\mathcal{F}_0 = \{[0,1]\}.$$

▶ Building \mathcal{F}_n : split $\begin{bmatrix} a \\ b \end{bmatrix}$, $\begin{bmatrix} c \\ d \end{bmatrix}$ ∈ \mathcal{F}_{n-1} , if $b + d \le n + 1$. Stern-Brocot partition SB_n :

- Base case: $SB_0 = \{[0, 1]\}.$
- ▶ Building SB_n : split $\left[\frac{a}{b}, \frac{c}{d}\right] \in SB_{n-1}$ always.



Farey partition

Farey partition \mathcal{F}_n :

• Base case: $\mathcal{F}_0 = \{[0,1]\}.$

▶ Building \mathcal{F}_n : split $\left[\frac{a}{b}, \frac{c}{d}\right] \in \mathcal{F}_{n-1}$ at mediant $\frac{a+c}{b+d}$, if $b+d \leq n+1$.



Properties:

- \mathcal{F}_k determines char. Sturmian word up to (k-1)-th symbol.
- The end-points \mathcal{F}_k are exactly $\{\frac{a}{b} \in \mathbb{Q} : 0 \le a \le b \le k+1\}$.
- Small number: $\Theta(k^2)$ intervals in \mathcal{F}_k

Farey partition

Farey partition \mathcal{F}_n :

• Base case: $\mathcal{F}_0 = \{[0,1]\}.$

▶ Building \mathcal{F}_n : split $\left[\frac{a}{b}, \frac{c}{d}\right] \in \mathcal{F}_{n-1}$ at mediant $\frac{a+c}{b+d}$, if $b+d \leq n+1$.



Properties:

- \mathcal{F}_k determines char. Sturmian word up to (k-1)-th symbol.
- The end-points \mathcal{F}_k are exactly $\{\frac{a}{b} \in \mathbb{Q} : 0 \le a \le b \le k+1\}$.
- Small number: $\Theta(k^2)$ intervals in $\mathcal{F}_k \Rightarrow$ entropy 0.

Weight of the Farey partition

Proposition

Farey partition is log-balanced a.e. with weight-function $f(n) = 2 \log n$.

Weight of the Farey partition

Proposition

Farey partition is log-balanced a.e. with weight-function $f(n) = 2 \log n$.

Farey intervals have comparable size almost everywhere:

Lemma

For almost every x, for large $n \ge n_0(x)$

$$\frac{1}{n^2} \le \left| I_n^{\mathcal{F}}(x) \right| \le \frac{(\log n)(\log \log n)}{n^2}$$



Figure. Histogram of interval sizes for n = 20. $\frac{1}{20^2} = 0.0025$, $\frac{1}{20} = 0.05$.

Stern-Brocot partition

Stern-Brocot partition SB_n :

• Base case: $SB_0 = \{[0, 1]\}.$

▶ Building SB_n : split $\left[\frac{a}{b}, \frac{c}{d}\right] \in SB_{n-1}$ always.



Stern-Brocot partition

Stern-Brocot partition SB_n :

• Base case: $SB_0 = \{[0, 1]\}.$

• Building SB_n : split $\left[\frac{a}{b}, \frac{c}{d}\right] \in SB_{n-1}$ always.



Associated to binary encoding of continued fractions:

$$[a_1, a_2, \ldots] \mapsto [0^{a_1 - 1}, 1, 0^{a_2 - 1}, 1, \ldots]$$

which follows construction of CFs by mediants.

Weight of the Stern-Brocot partition

Proposition

Stern-Brocot is log-balanced in measure with weight-function

$$f_{\mathcal{SB}}(n) = \frac{\pi^2}{6} \frac{n}{\log n}$$

Weight of the Stern-Brocot partition

Proposition

Stern-Brocot is log-balanced in measure with weight-function

$$f_{\mathcal{SB}}(n) = \frac{\pi^2}{6} \frac{n}{\log n}$$

Proposition

Stern-Brocot system of partitions is not log-balanced almost everywhere.

Weight of the Stern-Brocot partition

Proposition

Stern-Brocot is log-balanced in measure with weight-function

$$f_{\mathcal{SB}}(n) = \frac{\pi^2}{6} \frac{n}{\log n}$$

Proposition

Stern-Brocot system of partitions is not log-balanced almost everywhere.

Proof sketch.

– Depth in Stern-Brocot strongly related to the growth to sum of partial quotients $\sum_{k=1}^m a_k(x)$.

- Sum behaves well in measure but erratic almost-everywhere.

Consequences for our sources of zero-entropy

Corollary 1 Let ${\cal P}$ with $h({\cal P})>0$ and ${\cal SB}$ be the Stern-Brocot partition, then

$$L_n(x; \mathcal{P}, \mathcal{SB}) \sim \frac{6 h(\mathcal{P})}{\pi^2} \times n \log n$$
,

in measure.

Consequences for our sources of zero-entropy Corollary 1

Let ${\mathcal P}$ with $h({\mathcal P})>0$ and ${\mathcal{SB}}$ be the Stern-Brocot partition, then

$$L_n(x; \mathcal{P}, \mathcal{SB}) \sim \frac{6 h(\mathcal{P})}{\pi^2} \times n \log n$$
,

in measure.

Proof.

Since
$$f_{\mathcal{P}}(n) = h(\mathcal{P}) \times n$$
 and $f_{\mathcal{SB}}(m) = \frac{\pi^2}{6} \frac{m}{\log m}$ in measure,

$$\frac{\pi^2}{6} \frac{L_n(x; \mathcal{P}, \mathcal{SB})}{\log L_n(x; \mathcal{P}, \mathcal{SB})} \sim h(\mathcal{P}) \times n \,.$$

Applying logs shows that $\log L_n(x; \mathcal{P}, \mathcal{SB}) \sim \log n$ too.

Consequences for our sources of zero-entropy

Corollary 1 Let ${\mathcal P}$ with $h({\mathcal P})>0$ and ${\mathcal {SB}}$ be the Stern-Brocot partition, then

$$L_n(x; \mathcal{P}, \mathcal{SB}) \sim \frac{6 h(\mathcal{P})}{\pi^2} \times n \log n$$
,

in measure.

Corollary 2

Let \mathcal{P} with $h(\mathcal{P}) > 0$ and \mathcal{F} be the Farey partition, then

$$\log L_n(x; \mathcal{P}, \mathcal{F}) \sim \frac{h(\mathcal{P})}{2} \times n$$

almost everywhere.

Consequences for our sources of zero-entropy

Corollary 1

Let ${\mathcal P}$ with $h({\mathcal P})>0$ and ${\mathcal{SB}}$ be the Stern-Brocot partition, then

$$L_n(x; \mathcal{P}, \mathcal{SB}) \sim \frac{6 h(\mathcal{P})}{\pi^2} \times n \log n$$
,

in measure.

Corollary 2

Let ${\mathcal P}$ with $h({\mathcal P})>0$ and ${\mathcal F}$ be the Farey partition, then

$$\log L_n(x; \mathcal{P}, \mathcal{F}) \sim \frac{h(\mathcal{P})}{2} \times n$$

almost everywhere.

Proof.

For the input $f_1(n) = h(\mathcal{P}) \times n$, for the output $f_2(m) = 2 \log m$.

Second order term: continued fractions to Farey

Second order term might be irregular: big variability in L_n

Proposition

The Lochs' index from continued fractions to Farey satisfies

$$2\log L_n(x; \mathcal{CF}, \mathcal{F}) = h(\mathcal{CF}) \times n + c Z_n(x) \cdot \sqrt{n} + O(1) ,$$

where c > 0 and $Z_n \Rightarrow N(0, 1)$.

Recall. $f_{\mathcal{CF}}(n) = h(\mathcal{CF}) \times n$, and $f_{\mathcal{F}}(m) = 2 \log m$.

Second order term: continued fractions to Farey

Second order term might be irregular: big variability in L_n

Proposition

The Lochs' index from continued fractions to Farey satisfies

$$2\log L_n(x; \mathcal{CF}, \mathcal{F}) = h(\mathcal{CF}) \times n + c Z_n(x) \cdot \sqrt{n} + O(1) ,$$

where c > 0 and $Z_n \Rightarrow N(0, 1)$.

Recall. $f_{\mathcal{CF}}(n) = h(\mathcal{CF}) \times n$, and $f_{\mathcal{F}}(m) = 2 \log m$.

Proof.

Find specific formula for L_n in this case, then use CLT for $\log q_k(x)$.

A "non-example": Farey to continued fractions

Recall. $f_1(n) = 2 \log n$ not valid weight function a.e. for input \mathcal{P}^1 .

A "non-example": Farey to continued fractions

Recall. $f_1(n) = 2 \log n$ not valid weight function a.e. for input \mathcal{P}^1 .

Proposition

For the Farey ${\mathcal F}$ and the Continued Fraction ${\mathcal C}{\mathcal F}$ systems of partitions:

$$\lim_{n \to \infty} \frac{L_n(x; \mathcal{F}, \mathcal{CF})}{\log n} = \frac{12 \log 2}{\pi^2},$$

for almost every x.

A "non-example": Farey to continued fractions

Recall. $f_1(n) = 2 \log n$ not valid weight function a.e. for input \mathcal{P}^1 .

Proposition

For the Farey ${\mathcal F}$ and the Continued Fraction ${\mathcal C}{\mathcal F}$ systems of partitions:

$$\lim_{n \to \infty} \frac{L_n(x; \mathcal{F}, \mathcal{CF})}{\log n} = \frac{12 \log 2}{\pi^2},$$

for almost every x.

- Follows from characterization of L_n for the given sources.
- Main Theorem only gives this limit in measure



1. Definitions: partitions, Lochs' and weight function

2. Statement of main result and discussion

3. Examples of natural zero entropy sources that have weight

4. Concepts for the proof of the main result

5. Conclusions

Recall: main result

Theorem (Berthé, Cesaratto, R., Safe, 2021+)

Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with a.e. weight functions f_1 and f_2 . Then, under certain technical conditions

$$\lim_{n \to \infty} \frac{f_2\left(L_n(x; \mathcal{P}^1, \mathcal{P}^2)\right)}{f_1(n)} = 1,$$

for a.e. x.

The conditions are:

•
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

• f_2 is non decreasing ;

•
$$f_2(m+1) - f_2(m) = o(f_2(m))$$
 as $m \to \infty$.

Recall: main result

Theorem (Berthé, Cesaratto, R., Safe, 2021+)

Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with a.e. weight functions f_1 and f_2 . Then, under certain technical conditions

$$\lim_{n \to \infty} \frac{f_2\left(L_n(x; \mathcal{P}^1, \mathcal{P}^2)\right)}{f_1(n)} = 1,$$

for a.e. x.

The conditions are:

•
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

• f_2 is non decreasing ;

▶
$$f_2(m+1) - f_2(m) = o(f_2(m))$$
 as $m \to \infty$.

Intuition: $m = L_n$ satisfies $|I_m^{\mathcal{P}^2}(x)| \approx |I_n^{\mathcal{P}^1}(x)|$ up to log-terms.

Recall: main result

Theorem (Berthé, Cesaratto, R., Safe, 2021+)

Consider systems of partitions \mathcal{P}^1 and \mathcal{P}^2 , with a.e. weight functions f_1 and f_2 . Then, under certain technical conditions

$$\lim_{n \to \infty} \frac{f_2\left(L_n(x; \mathcal{P}^1, \mathcal{P}^2)\right)}{f_1(n)} = 1,$$

for a.e. x.

The conditions are:

•
$$\sum e^{-\delta f_1(n)} < \infty$$
 for every $\delta > 0$;

• f_2 is non decreasing ;

▶
$$f_2(m+1) - f_2(m) = o(f_2(m))$$
 as $m \to \infty$.

Intuition: $m = L_n$ satisfies $|I_m^{\mathcal{P}^2}(x)| \approx |I_n^{\mathcal{P}^1}(x)|$ up to log-terms.

 \Rightarrow Formal proof separated into two parts: upper-limit and lower-limit.

Proof-sketch: upper-limit

Upper-limit requires almost no conditions at all:

Lemma

Let \mathcal{P}^1 and \mathcal{P}^2 be a.e. log-balanced with weights f_1 and f_2 respectively. If f_2 is non-decreasing

$$\limsup_{n \to \infty} \frac{f_2(L_n(x; \mathcal{P}^1, \mathcal{P}^2))}{f_1(n)} \le 1 \qquad a.e.$$
Proof-sketch: upper-limit

Upper-limit requires almost no conditions at all:

Lemma

Let \mathcal{P}^1 and \mathcal{P}^2 be a.e. log-balanced with weights f_1 and f_2 respectively. If f_2 is non-decreasing

$$\limsup_{n \to \infty} \frac{f_2(L_n(x; \mathcal{P}^1, \mathcal{P}^2))}{f_1(n)} \le 1 \qquad a.e.$$

Proof-sketch for f_2 strictly increasing.

Fix $\varepsilon > 0$. Consider $m > f_2^{-1} \left((1 + \varepsilon) \times f_1(n) \right)$, then

$$-\log |I_m^{\mathcal{P}^2}(x)| \sim f_2(m) \ge (1+\varepsilon) \times f_1(n),$$

while $-\log |I_n^{\mathcal{P}^1}(x)| \sim f_1(n)$.

Proof-sketch: upper-limit

Upper-limit requires almost no conditions at all:

Lemma

Let \mathcal{P}^1 and \mathcal{P}^2 be a.e. log-balanced with weights f_1 and f_2 respectively. If f_2 is non-decreasing

$$\limsup_{n \to \infty} \frac{f_2(L_n(x; \mathcal{P}^1, \mathcal{P}^2))}{f_1(n)} \le 1 \qquad a.e.$$

Proof-sketch for f_2 strictly increasing.

Fix $\varepsilon > 0$. Consider $m > f_2^{-1} \left((1 + \varepsilon) \times f_1(n) \right)$, then

$$-\log |I_m^{\mathcal{P}^2}(x)| \sim f_2(m) \ge (1+\varepsilon) \times f_1(n),$$

while $-\log |I_n^{\mathcal{P}^1}(x)| \sim f_1(n)$. Thus $I_n^{\mathcal{P}^1}(x)$ too large for $I_m^{\mathcal{P}^2}(x)$.

Corollary

Let \mathcal{P} be a.e. log-balanced with weight f. If f is non-decreasing

$$\lim_{n \to \infty} \frac{f(L_n(x; \mathcal{P}, \mathcal{P}))}{f(n)} = 1 \qquad a.e.$$

Corollary

Let $\mathcal P$ be a.e. log-balanced with weight f. If f is non-decreasing

$$\lim_{n \to \infty} \frac{f(L_n(x; \mathcal{P}, \mathcal{P}))}{f(n)} = 1 \qquad a.e.$$

For zero-entropy sources, such as Sturmian words (Farey partition):

– knowing n digits allows us to deduce $L_n(x; \mathcal{P}, \mathcal{P}) \ge n$,

Corollary

Let $\mathcal P$ be a.e. log-balanced with weight f. If f is non-decreasing

$$\lim_{n \to \infty} \frac{f(L_n(x; \mathcal{P}, \mathcal{P}))}{f(n)} = 1 \qquad a.e.$$

For zero-entropy sources, such as Sturmian words (Farey partition):

- knowing n digits allows us to deduce $L_n(x; \mathcal{P}, \mathcal{P}) \ge n$,
- equality need not hold ! maybe $I_{n+1}(x) = I_n(x)$

Corollary

Let $\mathcal P$ be a.e. log-balanced with weight f. If f is non-decreasing

$$\lim_{n \to \infty} \frac{f(L_n(x; \mathcal{P}, \mathcal{P}))}{f(n)} = 1 \qquad a.e.$$

For zero-entropy sources, such as Sturmian words (Farey partition):

- knowing n digits allows us to deduce $L_n(x; \mathcal{P}, \mathcal{P}) \ge n$,
- equality need not hold ! maybe $I_{n+1}(x) = I_n(x)$
- weight function limits number of *digits* deduced.

Corollary

Let $\mathcal P$ be a.e. log-balanced with weight f. If f is non-decreasing

$$\lim_{n \to \infty} \frac{f(L_n(x; \mathcal{P}, \mathcal{P}))}{f(n)} = 1 \qquad a.e.$$

For zero-entropy sources, such as Sturmian words (Farey partition):

- knowing n digits allows us to deduce $L_n(x; \mathcal{P}, \mathcal{P}) \ge n$,
- equality need not hold ! maybe $I_{n+1}(x) = I_n(x)$
- weight function limits number of *digits* deduced.

Proof.

We know the upper-limit works. Lower-limit follows from

$$n \leq L_n(x; \mathcal{P}, \mathcal{P})$$
.

Proof-sketch: lower-limit

Lower-limit requires all of the conditions:

Lemma

Let \mathcal{P}^1 and \mathcal{P}^2 be a.e. log-balanced with weights f_1 and f_2 respectively, satisfying the conditions in the statement of the Theorem, then

$$1 \le \liminf_{n \to \infty} \frac{f_2(L_n(x; \mathcal{P}^1, \mathcal{P}^2))}{f_1(n)} \qquad a.e.$$

Proof-sketch: lower-limit

Lower-limit requires all of the conditions:

Lemma

Let \mathcal{P}^1 and \mathcal{P}^2 be a.e. log-balanced with weights f_1 and f_2 respectively, satisfying the conditions in the statement of the Theorem, then

$$1 \le \liminf_{n \to \infty} \frac{f_2(L_n(x; \mathcal{P}^1, \mathcal{P}^2))}{f_1(n)} \qquad a.e.$$

Proof techniques: covering argument + Borel-Cantelli.



1. Definitions: partitions, Lochs' and weight function

2. Statement of main result and discussion

3. Examples of natural zero entropy sources that have weight

4. Concepts for the proof of the main result

5. Conclusions

We have introduced the log-balance partitions \mathcal{P} with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)$, a.e. or in measure

We have introduced the log-balance partitions \mathcal{P} with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)\,, \quad \text{a.e. or in measure}$

 $\circledast \text{ Weight function intervenes naturally in change of basis} \\ \Rightarrow \text{ adapted } renormalization \text{ of the depths.}}$

We have introduced the log-balance partitions \mathcal{P} with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)$, a.e. or in measure

 $\circledast \text{ Weight function intervenes naturally in change of basis} \\ \Rightarrow \text{ adapted renormalization of the depths.}$

❀ Our results now apply to sources with zero or infinite entropy.

We have introduced the log-balance partitions \mathcal{P} with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)$, a.e. or in measure

- $\label{eq:Weight function intervenes naturally in change of basis} \Rightarrow adapted$ *renormalization*of the depths.
- $\circledast~\mbox{Our results}$ now apply to sources with zero or infinite entropy.
- \circledast We discussed zero-entropy sources from Number Theory

 \Rightarrow log-balanced, almost everywhere or just in measure.

We have introduced the log-balance partitions $\mathcal P$ with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)$, a.e. or in measure

- $\circledast \text{ Weight function intervenes naturally in change of basis} \\ \Rightarrow \text{ adapted } renormalization \text{ of the depths.}}$
- $\circledast~\mbox{Our results}$ now apply to sources with zero or infinite entropy.

Questions and further work

1. Obtain a general existence result for the weight ?

We have introduced the log-balance partitions $\mathcal P$ with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)$, a.e. or in measure

- $\circledast \text{ Weight function intervenes naturally in change of basis} \\ \Rightarrow \text{ adapted renormalization of the depths.}$
- $\circledast~\mbox{Our results}$ now apply to sources with zero or infinite entropy.

Questions and further work

- 1. Obtain a general existence result for the weight ?
- 2. Are the hypotheses necessary ?

We have introduced the log-balance partitions $\mathcal P$ with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)$, a.e. or in measure

- $\circledast \text{ Weight function intervenes naturally in change of basis} \\ \Rightarrow \text{ adapted renormalization of the depths.}$
- $\circledast~\mbox{Our results}$ now apply to sources with zero or infinite entropy.

Questions and further work

- 1. Obtain a general existence result for the weight ?
- 2. Are the hypotheses necessary ? \Rightarrow Limit applies for $\mathcal{F} \rightarrow C\mathcal{F}$ even though $f_{\mathcal{F}}(n) = 2\log n$

We have introduced the log-balance partitions $\mathcal P$ with weight f

 $-\log |I_n^{\mathcal{P}}(x)| \sim f(n)$, a.e. or in measure

 $\circledast \text{ Weight function intervenes naturally in change of basis} \\ \Rightarrow \text{ adapted renormalization of the depths.}$

- $\circledast~$ Our results now apply to sources with zero or infinite entropy.

Questions and further work

- 1. Obtain a general existence result for the weight ?
- 2. Are the hypotheses necessary ? \Rightarrow Limit applies for $\mathcal{F} \rightarrow C\mathcal{F}$ even though $f_{\mathcal{F}}(n) = 2\log n$
- 3. Results on average ?

Thank you!

References



A. Khintchine.

Metrische kettenbruchprobleme, Compositio Mathematica1, pp. 361–382, 1935.

G. Lochs,

Die ersten 968 Kettenbruchnenner von π , Monatsh. Math. 67, pp. 311–316, 1963.



C. Faivre.

A central limit theorem related to decimal and continued fraction expansion. Arch. Math. (Basel) 70, no. 6, pp 455-463, 1998.

K. Dajani, and A. Fieldsteel,

Equipartition of Interval Partitions and an Application to Number Theory,

Proceedings of the American Mathematical Society, vol 129, n. 12, pp. 3453-3460, 2001.